




StarmasterTM Pre-Show Lamb



Textured feed designed to fed to lambs weighing 70-140 pounds for that extra bloom prior to show.

Contains Decoquinate	Aids in prevention of Coccidiosis (see tag)
Contains cooked steam-flaked grains	Improves palatability, energy digestion and body condition.
Highly fortified	Promotes optimum performance.
Highly palatable	Encourages consistent consumption.
Nutritionally balanced and fortified with vitamins and minerals.	Promotes proper growth, development, reproduction and overall health.

DESCRIPTION: StarmasterTM Pre-Show Lamb feed is designed to follow StarmasterTM Lamb Stage 1 and continue the performance pace the lambs have begun. This high performance feed is formulated to achieve lean muscle growth, optimum body condition, muscle tone, and freshness in lambs weighing over 100 lbs. Provide clean, fresh water at all times.

FORM

Textured

BAG SIZE

50#



Starmaster™ Pre-Show Lamb

MEDICATED

(FOR LAMBS WEIGHING 70 TO 140 LB)

For the prevention of coccidiosis in young sheep caused by *eimeria bakuensis*, *eimeria crandallis*, *eimeria ovinoidalis* and *eimeria parva*.

ACTIVE DRUG INGREDIENT

DECOQUINATE13.6 G/TON

GUARANTEED ANALYSIS

CRUDE PROTEIN MIN 17.00 %
(THIS INCLUDES NOT MORE THAN 0.35% EQUIVALENT PROTEIN FROM NON- PROTEIN NITROGEN)
CRUDE FAT MIN 4.00 %
CRUDE FIBER MAX 15.00 %
CALCIUM MIN 1.75 % MAX 2.25 %
PHOSPHORUS MIN 0.70 %
SALT MIN 1.00 % MAX 1.50 %
SELENIUM MIN 0.30 PPM
VITAMIN A MIN 13,000 IU/LB

INGREDIENTS

GRAIN PRODUCTS, PLANT PROTEIN PRODUCTS, ANIMAL PROTEIN PRODUCTS, PROCESSED GRAIN BY-PRODUCTS, FORAGE PRODUCTS, ROUGHAGE, CANE MOLASSES, VEGETABLE OIL, CALCIUM CARBONATE, DICALCIUM PHOSPHATE, SALT, AMMONIUM CHLORIDE, SULFUR, FERROUS SULFATE, FERROUS CARBONATE, MANGANOUS OXIDE, MANGANESE SULFATE, ZINC OXIDE, ZINC SULFATE, COBALT CARBONATE, CALCIUM IODATE, SODIUM SELENITE, VITAMIN A SUPPLEMENT, VITAMIN D₃ SUPPLEMENT, VITAMIN E SUPPLEMENT, ANISE OIL.

DIRECTIONS FOR USE: FEED TO LAMBS AT THE RATE OF 1 POUND PER 30 POUNDS OF BODYWEIGHT TO PROVIDE DECOQUINATE AT THE RATE OF 22.7 MG PER 100 POUNDS OF BODYWEIGHT. FEED AT LEAST 28 DAYS DURING PERIODS OF EXPOSURE TO COCCIDIOSIS OR WHEN EXPERIENCE INDICATES THAT IT IS LIKELY TO BE A HAZARD. MIX THOROUGHLY WITH ANY ADDITIONAL GRAIN AND (OR) FORAGE BEING FED.

WARNING: DO NOT FEED TO SHEEP PRODUCING MILK FOR FOOD.

CAUTION: DO NOT USE IN FEEDS CONTAINING BENTONITE.

REMINDER: FEED IS PERISHABLE. STORE THIS PRODUCT IN A COOL, DRY AREA AWAY FROM RODENTS AND INSECTS.

WARNING: DO NOT OFFER ANY FEED THAT IS SPOILED, MOLDY, RODENT OR INSECT-INFESTED, OR ABNORMAL IN APPEARANCE OR ODOR, AS IT MAY CAUSE ILLNESS OR DEATH.

7922

Manufactured for: STARMaster FEEDS, P.O. BOX 789, THORNVILLE, OHIO